Unintentional Injury

Disease Information Packets – Slide Sets



Public Health Services,

Community Health Statistics

12/2012



What is Unintentional Injury?

- Unintentional injuries are injuries that are not caused on purpose and are free from harmful intent.
- Most unintentional injuries are related to:
 - Falls, poisonings/overdoses, motor vehicle crashes, struck by/against events, fires/burns, cuts/piercing, drowning/submersion, and overexertion.
- Unintentional injury is the 5th leading cause of death in the U.S.





Demographic Risk Factors

Age

 Those aged 65 years and older have the highest unintentional injury death and hospitalization rates compared to any other age group.

Ethnicity/race

 American Indians/Alaska Natives are more likely to die from unintentional injury compared to any other racial or ethnic group.

Gender

 Males are twice as likely to die from unintentional injury compared to females.

Socioeconomic status

 Those at a lower socioeconomic status may be at increased risk for unintentional injury.



Behavioral Risk Factors

Alcohol and drug abuse

- Alcohol use can lead to impaired brain function resulting in poor judgment, reduced reaction time, and loss of balance and motor skills, increasing the risk of unintentional injury.
- 1 in every 3 trafficrelated deaths involves a drunk driver.

- In 2010, alcohol-impaired drivers were involved in over 10,000 trafficrelated deaths.
- Drugs are involved in 18% of motor vehicle driver deaths.



Social and Behavioral Risk Factors

Lack of physical activity

- Lack of physical activity can lead to weak legs and decreased balance, which can increase the risk of falls for older adults.
- Weakened muscles from lack of physical activity can increase the risk of various mechanisms of unintentional injury (i.e., overexertion, falls).





National Statistics


- Unintentional is the leading cause of death for people aged 1-44.
- It is the 5th leading cause of death for all ages.
- In 2010, unintentional injury resulted in an estimated 120,000 deaths, 1.7 million hospitalizations, and 26 million ED discharges.

Top 10 Leading Causes of Death in the United States, 2009

- Heart Disease
- 2. Cancer
- 3. Respiratory Disease
- 4. Cerebrovascular Disease (Stroke)
- 5. Unintentional Injuries
- 6. Alzheimer' Disease
- 7. Diabetes
- 8. Influenza and Pneumonia
- 9. Nephritis
- 10. Suicide




National Statistics



• Falls were the most common cause of unintentional injury, followed by struck by/against injuries, overexertion, and motor vehicle-related injuries in 2009.




National Disparities



 Those aged 65 years and older are more likely to die or be hospitalized due to unintentional injury than any other age group.



National Disparities




 American Indians/Alaska Natives had the highest age-adjusted death rate for unintentional injury, whereas Asians/Pacific Islanders had the lowest in 2009.




Costs


- In 2009, unintentional injury cost the United States an estimated \$341.1 billion:
 - \$63.9 billion in direct medical costs
 - \$277.1 billion in lost productivity





















^{**}County actual rates are per 100,000 individuals. County age-adjusted rates per 100,000 2000 US standard population.

Source: Death Statistical Master Files (CADPH), CoSD, HHSA, Epidemiology & Immunization Services Branch; SANDAG, Current Population Estimates, 10/2010 Prepared by County of San Diego (CoSD). Health & Human Services Agency (HHSA). Community Health Statistics. 12/11/2012.



Prevention Strategies

Falls

- For children:
 - Supervising children when at play, installing stair gates, securing furniture to the wall, ensuring that children wear protective gear (i.e., helmets, wrist guards, knee/elbow pads) when playing active sports.
- For older adults:
 - Exercising regularly, wearing appropriate footwear (flat shoes with non-slip, thin soles), making the home safer by installing grab bars and non-skid mats in bathrooms, improving lighting, keeping frequently used items within easy reach, and removing loose rugs, electrical cords, and clutter in walkways.

Overdoses/Poisonings

- Locking medications, cleaning products, pesticides, and other household chemicals in their original containers and keep them out of reach.
- Never taking medicine in front of children or refer to is as candy.
- Reading labels on medications to insure proper dosage and avoid drug interactions.
- Seeking help for substance abuse problems.



Prevention Strategies

- Motor-Vehicle Related Injuries
 - Avoiding drinking and driving.
 - Enforcing seat belt laws and the use of booster seats for children under 4'9" in height.
 - Avoiding distractions while driving, such as talking or texting on cell phones, eating, or playing with the radio.
 - Obeying speed and traffic laws.
 - Enforcing graduated driver's license laws for new drivers.
- Struck By/Against Injuries
 - Wearing appropriate sports-related protective gear when engaged in sports-related activities.
 - Providing children safe places to play.
 - Remaining alert and aware of the surrounding environment.





Prevention Strategies

Overexertion Injuries

- Exercising regularly to strengthen muscles.
- Taking breaks during strenuous activities.
- Taking a firm stance and lift with your knees, keeping objects close to the body.
- Avoiding bending, reaching, and twisting when lifting.
- Getting help to move heavy or large objects.

• Cut/Pierce Injuries

- Keeping your eyes on your work while cutting.
- Using caution and wearing personal protective equipment when necessary.
- Using the right tool for the right job.



Contact Us

County of San Diego
Health and Human Services Agency
Public Health Services
Community Health Statistics
(619)285-6479

www.SDHealthStatistics.com

